

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : <b>A61K 31/55, C07D 495/04</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 97/23221</b> (43) International Publication Date: 3 July 1997 (03.07.97)</p>
<p>(21) International Application Number: PCT/US96/20048 (22) International Filing Date: 16 December 1996 (16.12.96) (30) Priority Data: 60/008,987 21 December 1995 (21.12.95) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): TRAN, Pierre, V. [US/US]; 11750 Forest Park Lane, Carmel, IN 46033 (US). (74) Agents: VORNDRAN-JONES, MaCharri et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).</p>		<p>(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report.</i></p>
<p>(54) Title: METHOD FOR TREATING DERMATITIS</p> <p>(57) Abstract</p> <p>The invention provides a method for treating fungal dermatitis comprising administering an effective amount of 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine to a patient in need thereof.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## METHOD FOR TREATING DERMATITIS

5           This invention relates to a method for treating  
fungal dermatitis using 2-methyl-4-(4-methyl-1-piperazinyl)-  
10H-thieno[2,3-b][1,5]benzodiazepine.

          Fungal dermatitis is a troublesome condition which  
may produce mild inflammation; wherein the organism producing  
10   the dermatitis may persist indefinitely, causing intermittent  
remissions and exacerbations of a gradually extending lesion  
with a scaling, slightly raised border. Often the condition  
results in a sudden visicular and bullous disease of the feet  
or an inflamed boggy lesion of the scalp.

15           It is known that fungal dermatitis may be caused by  
a variety of organisms. Most preferably, 2-Methyl-4-(4-  
methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine is  
used for the treatment of a condition selected from the group  
consisting of Tinea corporis, Tinea pedis, Tinea unguium,  
20   Tinea capitis, Tinea cruris, and Tinea barbae. It is  
preferred that the condition is selected from Tinea corporis  
and Tinea pedis. Most such infections respond to topical  
antifungal preparations such as the imidazoles. Certain  
cases may be resistant to such treatment and require systemic  
25   treatment using agents such as Griseofulvin; however, this  
agent is frequently associated with headaches,  
gastrointestinal distress, photosensitivity, rashes, or  
leukopenia. Angioedema has been reported, vertigo, and  
sometimes transient hearing reduction. Ketoconazole is an  
30   oral imidazole derivative which may be used for systemic  
treatment; however, this compound is associated with liver  
toxicity and other serious side effects. Therefore, effective  
agents having a favorable side effect profile are desired.  
Surprisingly, Applicant has discovered that 2-methyl-4-(4-  
35   methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine  
can be useful for treating fungal dermatitis.


-2-

The 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine compound is described in U.S. Patent No. 5,229,382 ('382), herein incorporated by reference in its entirety.

5 Fungal dermatitis can be particular troublesome to treat in non-compliant patients such as those suffering from psychotic conditions. The 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine compound can be particularly useful for treating fungal dermatitis in  
10 psychotic patients as the 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine compound can be administered as part of the patients' usual treatment regimen.

15 The presently claimed invention provides a method for treating fungal dermatitis comprising administering an effective amount of a 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine compound to a patient in need thereof.

20 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine, is a compound of Formula(I):



25 and is described in the '382 patent. The '382 patent teaches that 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-

b][1,5]benzodiazepine can be useful for the treatment of psychotic conditions and mild anxiety states.

For example, 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine has been assessed in a number of standard behavior tests predictive of antipsychotic activity. The claimed compound antagonized apomorphine-induced climbing behavior and hypothermia in mice. See Moore, N.A. et al Psychopharmacology 94 (2), 263-266 (1988). The compound also inhibits conditioned avoidance response in rats, and has been found to have a favorable profile of activity in a number of *in vitro* binding assays, designed to measure the degree of binding to neural receptors. For example, the compound is active at the dopamine D-1 and D-2 receptors as indicated by an IC50 of less than 1 uM in the 3H-SCH23390 (Billard, et al. *Life Sciences* 35:1885 (1984)) and the 3H spiperone (Seeman et al *Nature* 216:717 (1976)) binding assays respectively.

Further, 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine is active at the 5-HT<sub>2</sub> receptor and 5-HT<sub>1C</sub> receptor. The *in vitro* results would indicate that the compound is effective in the treatment of psychotic conditions but less likely to induce extra pyramidal side-effects.

To Applicant's knowledge none of these mechanisms of action are thought to be associated with the treatment, prevention, or amelioration of fungal dermatitis.

Surprisingly, and in accordance with the present invention, Applicant has discovered that 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine can be useful for treating or preventing fungal dermatitis.

As used herein the term "psychotic condition" shall refer to pathologic psychological conditions which are psychoses or may be associated with psychotic features. Such conditions include, but are not limited to the psychotic disorders which have been characterized in the DSM-III-R. Diagnostic and Statistical Manual of Mental Disorders, Revised, 3rd Ed. (1980). The DSM-III-R was

prepared by the Task Force on Nomenclature and Statistics of the American Psychiatric Association, and provides clear descriptions of diagnostic categories. The skilled artisan will recognize that there are alternative nomenclatures, nosologies, and classification systems for pathologic psychological conditions and that these systems evolve with medical scientific progress.

The usefulness of 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine for treating fungal dermatitis can be demonstrated by clinical trial.

Such effectiveness for the treatment, amelioration of fungal dermatitis was shown in the following clinical trial:

The study was a double-blind, randomized, parallel, placebo and haloperidol-controlled study of 55 human subjects. The study was conducted in 23 study centers throughout the United States and Canada. Patients were randomized to one of the five treatment groups: placebo, 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine 5 +/-2.5 mg/day, 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine 10 +/-2.5 mg/day, 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine 15 +/- 2.5 mg/day, or haloperidol (15 +/- 5 mg/day). The acute phase lasted for 6 weeks with evaluations performed weekly followed by a double-blind extension phase of at least one year.

Fungal dermatitis occurred statistically significantly less frequently in the 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine 15 +/- 2.5 mg/day than in the haloperidol in the double-blind extension treatment phase.

2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine is effective over a wide dosage range, the actual dose administered being dependent on the condition being treated. For example, in the

treatment of adult humans, dosages of from about 5 to 20 mg per day may be used. A once a day dosage is normally sufficient, although divided doses may be administered. For treatment of fungal dermatitis in a human, a dose range of from about 2.5 to 17.5 mg, preferably 5 to 20 mg per day is suitable. It is particularly preferred to administer about 15 to about 20 mg/day of 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine for the treatment of fungal dermatitis in a human. Radiolabelled 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine, can be detected in the saliva and thus the compound can potentially be monitored in patients to assess compliance.

The 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine compound will normally be administered orally for the treatment of fungal dermatitis, or may be administered by injection and, for this purpose, it is usually employed in the form of a pharmaceutical composition. Other suitable formulations are taught in the '382 patent. For the treatment of fungal dermatitis, a powder or transdermal formulation may be particularly desired.

As used herein, the term "mammal" shall refer to the Mammalia class of higher vertebrates. The term "mammal" includes, but is not limited to, a human. The term "treating" as used herein includes prophylaxis of the named condition or amelioration or elimination of the condition once it has been established.


For the treatment of fungal dermatitis, the patient may be a non-human mammal. In such instances, the 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine compound may be administered as a feed additive, tablet, or transdermally.

Methods for preparing 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine are provided by the '382 patent; however, the following examples may be instructive as well.

Example 1

Technical Grade 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine

5



Intermediate 1

In a suitable three neck flask the following was added:

10

Dimethylsulfoxide (analytical): 6 volumes

Intermediate 1 : 75 g

N-Methylpiperazine (reagent) : 6 equivalents

15 Intermediate 1 can be prepared using methods known to the skilled artisan. For example, the preparation of the Intermediate 1 is taught in the '382 patent.

20 A sub-surface nitrogen sparge line was added to remove the ammonia formed during the reaction. The reaction was heated to 120°C and maintained throughout the duration of the reaction. The reactions were followed by HPLC until  $\leq 5\%$  of the intermediate 1 was left unreacted. After the reaction was complete, the mixture was allowed to cool slowly to 20°C (about 2 hours). Each reaction mixture was then transferred to an appropriate three neck round bottom flask and water bath. To this solution with agitation was added 10 volumes reagent grade methanol and the reaction was stirred at 20°C for 30 minutes. Three volumes of water was added slowly over

25



-7-

about 30 minutes. The reaction slurry was cooled to zero to 5°C and stirred for 30 minutes. The product was filtered and the wet cake was washed with chilled methanol. The wet cake was dried in vacuo at 45°C overnight. The product was  
5 identified as technical 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine..

Yield: 76.7%; Potency: 98.1%

The procedure of Example 1 was repeated substantially as  
10 described above and provided a yield of 81% with a potency of 101.1%.

#### Example 2

15 Technical Grade 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine

Intermediate 1 (*supra*) was suspended in DMSO (3.2 vol.) and toluene (4.5 vol.). A portion (=0.65 vol.) of the solvent was removed by distillation at 120-125°C. The  
20 mixture was cooled to 110°C, N-methylpiperazine(NMP, 4.2 equiv.) was added and the mixture heated back to reflux (120-125°C). Another portion (=1 vol.) of the solvent was removed by distillation to dry the reaction mixture. A vigorous reflux was desired to drive the reaction to completion (about  
25 7 hrs.) by removing ammonia from the reaction. The product was isolated by the slow addition of water (12.75 vol.) to the cooled (10°C) reaction solution. The product was collected by filtration and washed with chilled water (2 vol.). The crude 2-methyl-4-(4-methyl-1-piperazinyl)-10H-  
30 thieno[2,3-b][1,5]benzodiazepine was dried in vacuo at 60°C. The product was recrystallized from hot toluene (5 vol.) to give a technical grade 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine. After drying in vacuo at 50°C, the technical grade 2-methyl-4-(4-methyl-1-  
35 piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine was recrystallized again from ethyl acetate (10 vol.)/toluene (0.62 vol.)/methanol (3.1 vol.) to give 2-methyl-4-(4-methyl-

1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine as a methanol solvate. The methanol solvate upon drying at >50°C was converted to an anhydrous technical grade 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine.

5

### Example 3

#### Tablet Formulation

A portion of the hydroxypropyl cellulose was dissolved in purified water to form a solution for granulation. The remaining hydroxypropyl cellulose (total of 4.0% w/w final tablet weight), which was an extra fine grade, was combined with the 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine compound (1.18% w/w), lactose (79.32% w/w) and a portion of the crospovidone (5% w/w) in a high shear granulator. All ingredients were security sieved prior to addition and dry blended in the granulator. This mixture was then granulated with the hydroxypropyl cellulose solution in the high shear granulator. The granulation was wet sized using standard methods. The wet granulation was then dried in a fluidized bed dryer and sized. The material was then added to a tumble bin mixer.

The outside powders consisting of microcrystalline cellulose (granular) (10% w/w), magnesium stearate (0.5% w/w), and the remainder of the crospovidone were added to the sized granulation. The mixture was blended and compressed with the appropriate tooling on tablet compression equipment.

30

#### Subcoating:

Hydroxypropyl methylcellulose (1.5% w/w) was mixed with purified water to form a solution. Core tablets were divided into approximately equal sections and spray coated with the hydroxypropyl methylcellulose solution. The operation was performed in a perforated coating pan.

35

Coating of Core Tablets:

5           Color Mixture White (hydroxypropyl methylcellulose,  
polyethylene glycol, polysorbate 80, and titanium dioxide)  
was mixed with purified water to form the coating suspension.  
Subcoated tablets were divided into approximately equal  
sections and spray coated with the coating suspension  
described above. The operation was performed in a perforated  
10 coating pan.

The coated tablets were lightly dusted with  
carnauba wax and imprinted with appropriate identification.

We Claim:

1. A method for treating fungal dermatitis comprising administering an effective amount of 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine or a pharmaceutically acceptable salt thereof to a patient in need of such treatment.
2. A method of Claim 1 wherein the effective amount is from about 5 to about 20 mg/day for a human patient.
3. A method of Claim 2 wherein the effective amount is from about 15 mg/day to about 20 mg/day.
4. A method of Claim 3 wherein the 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine is administered as a tablet.
5. A method of Claim 3 wherein the effective amount is administered using a transdermal formulation.
6. A method of Claim 1 wherein the patient is a mammal.
7. A method of Claim 6 wherein the 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine is administered as a feed additive.
8. A method of Claim 6 wherein the patient is a human.
9. A method of Claim 8 wherein the patient is not diagnosed with a psychotic condition.
10. A method of Claim 1 wherein the patient suffers from fungal dermatitis.

11. A method of Claim 10 wherein the fungal dermatitis is selected from the group consisting of Tinea corporis, Tinea pedis, Tinea unguium, Tinea capitis, Tinea cruris, and Tinea barbae.

12. A method of Claim 11 wherein the fungal dermatitis is selected from the group consisting of Tinea corporis and Tinea pedis.

13. A method of Claim 8 wherein the patient is diagnosed with a psychotic condition.

14. 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine or a pharmaceutically acceptable salt thereof for treating fungal dermatitis.

15. The use of 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of fungal dermatitis.

16. A pharmaceutical formulation adapted for the treatment of fungal dermatitis containing 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/20048

**A. CLASSIFICATION OF SUBJECT MATTER**  
 IPC(6) :A61K, 31/55; C07D, 495/04  
 US CL :5H, 220; 540, 557  
 According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**  
 Minimum documentation searched (classification system followed by classification symbols)  
 U.S. : 514, 220; 540, 557

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 5,229,382 (CHAKRABARTI) 20 July 1993 (20.07.93).	1-16

Further documents are listed in the continuation of Box C.  See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A' document defining the general state of the art which is not considered to be of particular relevance	*X'	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B' earlier document published on or after the international filing date	*Y'	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L' document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z'	document member of the same patent family
*O' document referring to an oral disclosure, use, exhibition or other means		
*P' document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 17 MARCH 1997	Date of mailing of the international search report <b>08 APR 1997</b>
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>Robert T. Bond</i> ROBERT T. BOND aco Telephone No. (703) 308-1235